PLEASE CONSIDER SENDING YOUR PRESCRIPTION ELECTRONICALLY. ALL OF OUR PHARMACY LOCATIONS ACCEPT ELECTRONIC PRESCRIPTIONS. Note: This form is intended for prescriber use only, if faxed, the fax must come from MD office or hospital (may not be faxed by patient).



Walgreens Pharmacy		Prescription/Pharmacy Intake F	orm
Pharmacy:			
Pharmacy Fax:			
Date Needed: Ship To: □ Prescriber's Office □ Patient's Home	☐ Other:		
PATIENT INFORMATION			
Patient name:		_ DOB: □Male □ Femal	le
Address:			
City:	State:	Zip code:	
Phone # (Daytime):			
E-mail Address:	Case Manager:		
Insurance provider (Please include copy of front and back of card):			
ID #:Policy/Group #:		□ Patient is eligible for Medic	are
Name of Insured:	Employer:	2 1 2 1	
Relationship to Patient: Self Other:	Prescription Card: ☐ Yes ☐ No C	Jarrier: Policy/Group #:	
Will there be access to anaphylactic medications and oxygen at the administration site?			
CLINICAL ASSESSMENT - Please complete ALL sections to avo			
□ Patient is new to therapy □ Patient is restarting therapy □ Patient is currently			
Diagnosis: ☐ Moderate persistent asthma, uncomplicated (J45.40) ☐ Severe persistent asthma			
□Pulmonary eosinophilia, not elsewhere classified (J82) □Other		Date of Diagnosis:	
Other Diagnosis/Conditions:			
Eosinophil count Cells/µL IgE Level IU/mL Current Weight:			
Other Therapies Tried & Failed (Please List):			
Allergies:			
PRESCRIPTION INFORMATION			
□ Cinqair (reslizumab) 100 mg/10 mL vial	□ Nucala (mepolizumab)		
3 mg/kg once every 4 weeks by intravenous infusion over 20 to 50 minutes. To be administered by a healthcare professional.		ore-filled autoinjector □ 100 mg/mL pre-filled syringe □ 40 mg/mL	illed syringe
Quantity: Refills:	☐To be administered by a SEVERE ASTHMA	nearthcare professional.	
□ Dupixent (dupilumab)		usly once every 4 weeks into the upper arm, thigh or abdomen.	
\square 100 mg pre-filled syringe \square 200 mg pre-filled syringe \square 200 mg pre-filled pen		sly once every 4 weeks into the upper arm, thigh or abdomen.	
□300 mg pre-filled syringe □300 mg pre-filled pen		OMATOSIS WITH POLYAGNIITIS (EGPA)	
☐ Initial dose of 400 mg (two 200 mg injections at different injection sites) followed by		te 100 mg subcutaneous injections once every 4 weeks	
200 mg subcutaneously once every other week into the thigh or abdomen ☐ Initial dose of 600 mg (two 300 mg injections at different injection sites) followed by	into the upper arm, thig Quantity: Refills		
300 mg subcutaneously once every other week into the thigh or abdomen	☐ Tezspire (tezepelumab-ekko)		
☐ 100 mg subcutaneously once every other week into the thigh or abdomen	□210 mg vial □210 mg pre-1		
\square 200 mg subcutaneously once every other week into the thigh or abdomen		once every 4 weeks administered by a healthcare professional	
☐ 300 mg subcutaneously once every other week into the thigh or abdomen	Quantity: Refills		
☐ 300 mg subcutaneously once every 4 weeks into the thigh or abdomen	□Xolair (omalizumab) 150 mg v	/ial kit	
Quantity:Refills: □ Fasenra (benralizumab) 30 mg/mL pre-filled syringe	☐ Supply Kit (#2) 18q 1 & ½ syringe 3m	I	
30 mg administered once every 4 weeks for the first 3 doses, and then once every 8 weeks	(#2) 25g 5/8 safety needle		
thereafter by subcutaneous injection into the upper arm, thigh or abdomen.	(#2) alcohol swabs		
To be administered by a healthcare professional.	☐ Xolair PFS (omalizumab)		
Quantity:Refills: □ Fasenra (benralizumab) 30 mg/mL Pen		nge □150 mg/1 mL pre-filled syringe	
30 mg administered once every 4 weeks for the first 3 doses, then once every 8 weeks	☐ To be administered by a	a nealthcare professional. as patient received at least 3 doses of Xolair under	
thereafter by subcutaneous injection into the upper arm, thigh or abdomen.		as patient received at least 3 doses of Xolair under provider with no hypersensitivity reactions? ☐ Yes ☐ No	
Quantity:Refills:	Every 4 weeks dosing:	Every 2 weeks dosing:	
Other:	☐75 mg per dose subcutaned		•
Directions: Quantity:Refills:	☐ 150 mg per dose subcutane	, ,	•
Other:	□ 225 mg per dose subcutane		every 2 weeks.
Directions:	☐ 300 mg per dose subcutane Quantity: Refills		
Quantity:Refills:	Į.		
I authorize, by my signature below, the dispensing of appropriate needles and syringes, in		the administration of injectable products by patient or caregiver.	Authorization fo
supplies runs concurrently with the number of refills or time frame specified for the drug. PRESCRIBER INFORMATION	·		
Prescriber's name:	Practice/facility:		
Address:	Practice/facility: City:		
Office contact:		State:Zip code:	
Email:	best time to call:	Preferred method of contact: □Email □Phone □Fax	i
State license #:DEA #:	INPI #:	Medicald UPIN #:	cally possessy and
In order for a brand name product to be dispensed, the prescriber must handwrite "Brand Necessary" or "Bran that the information above is accurate to the best of my knowledge. Prescriber's signature required on one of the		unio required ranguage after their signature. I certify that the above therapy is medic	Jany necessary and
Dispense as written	Substitution permitted	Date	

*** THIS FORM IS NOT VALID IN THE STATE OF ALABAMA ***